arXiv:1512.00686v2 [math.GT] 10 Dec 2015

NEW INVARIANTS OF LINKS FROM A SKEIN INVARIANT OF COLORED LINKS

FRANCESCA AICARDI

Abstract. New invariants of links are constructed by using the skein invariant polynomial of colored links defined in [1]. These invariants are stronger than the homflypt polynomial.

Introduction

This note refers to the work [1], in which I have defined, via skein relation, an invariant of colored links. Let me recall some definitions.

An oriented link with n components is a set of n disjoint smooth oriented closed curves embedded in S^3. An oriented link is *colored*, if each component is provided with a color. In other words, a link is colored if a function γ is defined on the set of components to a finite set H of colors. Let's call *coloration* such a function γ. Every coloration introduces an equivalence relation in the set of components: two components are equivalent if they have the same color. Let $\mathbf{C}$ be the set of components of a link L. Two colorations γ and γ' of L are said *equivalent* if there is a bijection in H between the images $\gamma(\mathbf{C})$ and $\gamma'(\mathbf{C})$. Two colorations define the same partition of the set $\mathbf{C}$ into classes of equivalence if and only if they are equivalent.

An invariant of colored links takes the same value on isotopic links with the same coloration, and may take different values on the same link with different colorations. The invariant F of colored links defined in [1] takes the same value on isotopic links with equivalent colorations. In Section 1, the statement of the theorem defining the invariant F is reported.

The invariant F has three variables, say x, w and t. There are two natural ways to define an invariant of classical links by using F. The first way, is to consider a link as a colored link, whose components have all the same color. In this case the invariant F becomes independent of x, and is in fact another instance of the homflypt polynomial. In particular, it becomes the Jones polynomial in the variable t setting $w = t^{1/2}$. Given a classical link L, we will denote $F^1(L)$ the invariant F on the colored link obtained by coloring L with a sole color.

The second way, is to consider a link as a colored link, whose components have all different colors. In this case F yields an invariant which is stronger than the homflypt polynomial. Given a classical link L_n with n components, we will denote $F^n(L_n)$ the invariant F on the colored link obtained by coloring L_n with n colors.

Among the 89 pairs of non-isotopic links with at most 11 crossings, non distinguished by the homflypt polynomial, 49 pairs contain links with three components. I have computed F^3 on 20 pairs of such pairs: six of these pairs are distinguished by F^3 (see Proposition 3 in Section 3).

However, there are other ways to define an invariant of classical links by using F. Indeed, we can calculate F on the whole set of colored links obtained from a link L with n components by coloring it by all the possible non-equivalent colorations. Then, for every integer partition p of n in c parts ($c \leq n$), we take all partitions of the set of n components in c blocks, whose cardinalities correspond to the partition p. These partitions of the set of components are said

1991 *Mathematics Subject Classification.* 57M25 .

Key words and phrases. knot theory, knots invariants.

to be of type p. We thus define the invariant $\mathbf{F}_p^c$ as the set of values taken by F on the colored links obtained by coloring L with c colors according to each one of the type p partitions, i.e., by coloring the components with the same color if and only if they belong to the same block. Example: the link below has three components, that are colored according to the partitions indicated.



The invariant $\mathbf{F}^2_{(2,1)}$ distinguishes pairs of the above mentioned pairs of non isotopic links, which are not distinguished by F^3.

As we have said in [1], the definition of F is a byproduct of the study of another class of links, the *tied links* (see [2]). In fact, the tied links are links whose components may be connected in several points by ties. But these ties behave exactly as an equivalence relation between the link components, so that isotopic tied links are isotopic colored links with equivalent colorations (see Section 1 for more precise definitions). Tied links can be seen also as the closure of tied braids, that constitute a diagrammatic representation of the elements of an abstract algebra, the so-called nowadays bt-algebra (algebra of braids and ties). In [2] we have defined, via skein relation, a polynomial $\mathcal{F}$, invariant of tied links, and we have shown that $\mathcal{F}$ can be realized also by the Jones recipe, using the trace on the bt-algebra established in [3]. The properties of the trace allowed us to make a computer program to calculate $\mathcal{F}$. We may define a tied link TL from a colored link L so that $F(L)$ is recovered from $\mathcal{F}(TL)$ by a variable change. This has allowed me to verify the values of F obtained on colored links by the skein relation, presented in this paper.

Finally, I show some unexpected identities satisfied from the values forming the invariants F_p^c for the pairs distinguished by such invariants, that allows to formulate a conjecture. The calculation of F_p^c on all pairs of three-component links not distinguished by the homflypt polynomial has to be completed in order to understand these identities or to disprove the conjecture.

1. The invariant F of colored links

Definition 1. Two oriented colored links are *c-isotopic* if they are ambient isotopic, and their colorations are equivalent.

A colored link diagram is like the diagram of a link, where each component is colored.

Let R be a commutative ring and $\mathcal{L}_c$ be the set of colored oriented link diagrams. By *invariant of colored links* we mean a function $I : \mathcal{L}_c \to R$, which is constant on each class of c-isotopic links.

Remark 1. In the sequel, we denote by L an oriented colored link as well as its diagram, if there is no risk of confusion.

Theorem 1. *There exists a rational function in the variables x, w, t, $F : \mathcal{L}_c \to \mathbb{Q}(x, t, w)$, invariant of oriented colored links, uniquely defined by the following three conditions on colored-link diagrams:*

I *The value of F is equal to 1 on the unknotted circle.*

II *Let L be a colored link, with n components and c colors. By $[L; \mathsf{O}]$ we denote the colored link with $n+1$ components consisting of L and the unknotted and unlinked circle, colored with the $(c+1)$-th color. Then*

$$F([L; \mathsf{O}]) = \frac{1}{wx} F(L).$$

III *(skein relation) Let L_+ and L_- be the diagrams of two colored links, that are the same outside a small disc into which two strands enter, and inside this disc look as shown in figure 1a. Black and gray indicate any two colors, non necessarily distinct. Similarly, let $L_\sim$ and $L_{+,\sim}$ be two links that inside the disc look as shown in figure 1b, while outside the disc coincide with L_+ and L_-, except -possibly- for the colors of all the components of the link having the same colors as the strands entering the disc: all these components in $L_\sim$ and $L_{+,\sim}$ are colored by a sole color. Then the following identity holds:*

$$\frac{1}{w} F(L_+) - wF(L_-) = \left(1 - t^{-1}\right) F(L_\sim) + \frac{1}{w}(1 - t^{-1}) F(L_{+,\sim}).$$



FIGURE 1. a) The discs where L_+ and L_- are not c-isotopic. b) The discs where $L_\sim$, $L_{+,\sim}$, and $L_{-,\sim}$ are not c-isotopic. Blue and green indicate any two colors.

Remark 2. The skein relation III holds for *any* two colors of the involved strands. In particular, if the two colors coincide, then $L_+ = L_{+,\sim}$ and $L_- = L_{-,\sim}$, so that relation III is reduced to the relation

IV
$$\frac{1}{tw} F(L_{+,\sim}) - wF(L_{-,\sim}) = (1 - t^{-1}) F(L_\sim).$$

Relation III also implies following two relations; they will be used in the sequel.

Va
$$\frac{1}{w} F(L_+) = w\left[F(L_-) + (t-1)\, F(L_{-,\sim})\right] + (t-1)\, F(L_\sim).$$

Vb
$$wF(L_-) = \frac{1}{w}[F(L_+) + (t^{-1} - 1) F(L_{+,\sim})] + (t^{-1} - 1) F(L, \sim).$$

Remark 3. As an example of using relations I, II and III, let's calculate the value of F on the unlink O_n^c, consisting of n circles and c colors. If $n = c$, then, using II,

$$F(\mathsf{O}_c^c) = \frac{1}{wx} F(\mathsf{O}_{c-1}^{c-1}) = \left(\frac{1}{wx}\right)^{c-1}.$$

If $n > c$, then there are two circles colored by the same color. Regard them as belonging to $L_\sim$. Then $L_+ = L_- = \mathsf{O}_{n-1}^c$ (see figure).



By IV, $(1 - t^{-1})F(\mathsf{O}_n^c) = \left(\frac{1}{tw} - w\right)F(\mathsf{O}_{n-1}^c)$. So

$$F(\mathsf{O}_n^c) = \frac{1 - tw^2}{tw(1 - t^{-1})} F(\mathsf{O}_{n-1}^c) = \frac{tw^2 - 1}{(1 - t)w} F(\mathsf{O}_{n-1}^c) := \frac{y}{xw} F(\mathsf{O}_{n-1}^c),$$

where

$$y = \frac{x(tw^2 - 1)}{1 - t}. \tag{1}$$

By induction, $F(\mathsf{O}_n^c) = \left(\frac{y}{xw}\right)^{n-c} F(\mathsf{O}_c^c)$. Thus

$$F(\mathsf{O}_n^c) = \frac{y^{n-c}}{(wx)^{n-1}}. \tag{2}$$

Remark 4. Let L be a knot or a link whose components have all the same color. Then $F(L)$ is defined by I and IV. Recall that the homflypt polynomial P is defined by the condition that $P(\mathsf{O}) = 1$ and the skein relation

$$\ell P(L_{+,\sim}) + \ell^{-1} P(L_{-,\sim}) + m P(L_\sim) = 0, \tag{3}$$

where ℓ and m are two complex variables. Comparing the skein relation IV, multiplied by $t^{1/2}$, with the equation above, we see that F is a homflypt polynomial with

$$\ell = i\frac{1}{w\sqrt{t}} \quad \text{and} \quad m = i\left(\frac{1}{\sqrt{t}} - \sqrt{t}\right).$$

In particular, for $w = t^{1/2}$ it becomes the Jones polynomial.

Remark 5. The polynomial F is multiplicative with respect to the connected sum of colored links L_1 and L_2 in the following sense. The components C_1 and C_2 of the two links that merge must have the same color, while the colors of the components in L_1 not equal the color of C_1 must be different from the colors of the components in L_2 not equal to the color of C_2.

2. The new invariants of links

Let L be a classical link with n components. We denote by $\mathbf{C}_n := \{C_1, C_2, \ldots, C_n\}$ the ordered set of the components of L. A set-partition

$$\mathbf{I}_j := (I_1, I_2, \ldots, I_{c_j})$$

of $\mathbf{C}_n$, contains $c_j \leq n$ non void blocks. For every set-partition $\mathbf{I}_j$ of $\mathbf{C}_n$, we define a colored link L_j in this way: we color the components of L with c_j colors, in such a way that the components entering the same block have the same color. Observe that the number N of non-equivalent colorations of the link L satisfies $N \leq B_n$, where B_n is the n-th Bell number. The equality holds in general except in particular cases of symmetries among the components of the link.

Definition 2. Given a set partition $\mathbf{I} := (I_1, I_2, \ldots, I_c)$ of $\mathbf{C}_n$, let $n_i = |I_i|$, so that $n_1 + n_2 + \cdots + n_c = n$. We call the integer partition p of n in c part,

$$p := (n_1, n_2, \ldots, n_c) \qquad n_i \geq n_{i+1}$$

the *type* of the partition $\mathbf{I}$.

For instance, if $n = 4$, among the following partitions of $\mathbf{C}_4$ in two blocks:

$$\mathbf{I}_1 := (\{C_1, C_2\}, \{C_3, C_4\}), \quad \mathbf{I}_2 := (\{C_1\}, \{C_2, C_3, C_4\}), \quad \mathbf{I}_3 := (\{C_1, C_4\}, \{C_2, C_3\}),$$

the partitions $\mathbf{I}_1$ and $\mathbf{I}_3$ have the same type $(2, 2)$, and $\mathbf{I}_2$ has type $(3, 1)$.

Definition 3. For every partition p of n in c parts ($1 \leq c \leq n$), let N_p^c be the number of partitions of $\mathbf{C}_n$ of type p which define non-equivalent colorations of L with c colors, and $L_1, \ldots, L_{N_p^c}$ be the corresponding colored links. The invariant $\mathbf{F}_p^c$ of L is the non-ordered set

$$\mathbf{F}_p^c(L) := \{F(L_1), F(L_2), \ldots, F(L_{N_p^c})\}.$$

Remark 6. It is evident that for a link L wit n components, if $c = 1$, $p = (n)$, so $N_n^1 = 1$, and $\mathbf{F}_{(n)}^1(L) = \{F^1(L)\}$, i.e., the value of F on the link L with all component colored with the same color. Furthermore, for $c = n$ the invariant $\mathbf{F}^n_{\underbrace{(1,1,\ldots,1)}_{n}} = \{F^n(L)\}$.

Remark 7. In the following figures, two arrows going from a link L with a marked crossing, to two colored links L_1 and L_2, indicate that $F(L) = \alpha F(L_1) + \beta F(L_2)$, where α and β are given by relation IV (Remark 2); three arrows going from a colored link L with a marked crossing, to three links L_1, L_2 and L_3, indicate that $F(L) = \gamma F(L_1) + \delta F(L_2) + \lambda F(L_3)$, where γ, δ and λ are given by relations Va or Vb, depending on the sign of the marked crossing.

2.1. **An example.** We give here the experimental evidence that for $c > 1$, $\mathbf{F}_p^c$ is more powerful than the homflypt polynomial.

Consider, for instance, the links A and B, shown in 3. They have four components, six positive crossings, and are colored with a sole color. A and B are evidently non isotopic. The calculation of $F(A)$ and $F(A)$ shows that A and B are not distinguished by the homflypt polynomial (see Remark 4). Indeed, applying the skein relation IV to the crossings marked in Figure 2, we obtain

$$F(A) := F(A_{+,\sim}) = tw^2 F(A_{-,\sim}) + (t-1)wF(A_{\sim}),$$

$$F(B) := F(B_{+,\sim}) = tw^2 F(B-,\sim) + (t-1)wF(B_{\sim}).$$

But $A_{-,\sim} = B_{-,\sim} = L_1$ and $A_{\sim} = B_{\sim} = L_2$. Therefore F coincides on A and on B.



FIGURE 2. $F(A)$ and $F(B)$ coincide

Consider now the colored links obtained by coloring A and B with two colors, more precisely three components with one color and the remaining component by another color (i.e., corresponding to the partitions of type $p = (3, 1)$). There are two of such non-equivalent colorations of A and of B, giving the colored links A_1, A_2 and B_1, B_2 respectively (see Figure 3)



FIGURE 3. Calculation of $\mathbf{F}^2_{(3,1)}(A)$ and $\mathbf{F}^2_{(3,1)}(B)$

Proposition 1. *The links A and B are distinguished by the invariant $\mathbf{F}^2_{3,1}$.*

Proof. We have $\mathbf{F}^2_{(3,1)}(A) = \{F(A_1), F(A_2)\}$ and $\mathbf{F}^2_{(3,1)}(B) = \{F(B_1), F(B_2)\}$. From Figure 3 we see that $F(B_2) = F(A_2)$. We calculate now $F(A_1)$ and $F(B_1)$.

We have:
$F(A_1) = w^2tF(A_{11}) + w(t-1)F(A_{12})$,
$F(A_{11}) = F(A_{12})y/(wx)$, $F(A_{12}) = w^2F(A_{121}) + w^2(t-1)F(A_{122}) + w(t-1)F(A_{123})$,
$F(A_{121}) = F(A_{212})y/(wx)$, $F(A_{122}) = F(A_{123})y/(wx)$,
$F(A_{123}) = w^2tF(A_{1231} + w(t-1)F(A_{1232})$,
$F(A_{212}) = w^2F(A_{2121}) + w^2(t-1)F(A_{2122}) + w(t-1)F(A_{2123})$
$F(A_{2121}) = 1/(wx))$, $F(A_{1231} = F(A_{2122}) = y/(wx)$, $F(A_{1232}) = F(A_{2123}) = 1$.
We get
$F(A_1) = (1 - xt^5w^6 - t - tw^2 + t^2 - 2xtw^2 + 2xt + w^4t^2x - t^3w^2 + t^3w^4 + 2xt^2w^2 - 6xt^3w^2$
$- 3xt^2 + 4xt^3 + 2xt^3w^4 - xt^4w^6 + 3xt^4w^2 + 3xt^5w^4 - 3xt^5w^2 - 2xt^4 + xt^5)w^3/(x(t-1)^2)$.

On the other hand, we have
$F(B_1) = w^2F(B_{11}) + w^2(t-1)F(B_{12}) + w(t-1)F(B_{13})$,
$F(B_{11}) = F(A_{11})$, $F(B_{12}) = F(B_{13})y/(wx)$, $F(B_{13}) = w^2tF(B_{131}) + w(t-1)F(B_{132})$,
$F(B_{131}) = F(B_{132})y/(wx)$, $F(B_{132}) = F(A_{123})$.
Therefore,
$F(B_1) = w^3(w^4t^2 + 3w^4t^2x - 6xt^3w^2 - 3xt^2 + 4xt^3 - 2xt^4 - w^6t^3x - xt^4w^6 + 3xt^4w^2 - xt^5w^6$
$+ 3xt^5w^4 - 3xt^5w^2 + xt^5 + 3xt^3w^4 - 3xtw^2 + 3xt + 1 - 2tw^2)/(x(t-1)^2)$.

Since $F(B_1) \neq F(A_1)$, the links A and B are distinguished by $\mathbf{F}^2_{(3,1)}$. □

3. Non trivial examples

In this section we show some pairs of links which are not distinguished by the homflypt polynomial, but that are distinguished by $\mathbf{F}^c_p$.

The calculation of F by the skein relation is shown in the figures for some pairs. For the meaning of the arrows, see Remark 7. The skein-trees terminate in some simple basic colored links, belonging to a list $\mathsf{L1}, \mathsf{L2}, \ldots, \mathsf{L41}$, shown in Figure 12. The values of F on these colored links are given in Appendix 1.

The links are named according to LinkInfo, see [4].

Proposition 2. *The invariant $\mathbf{F}^3_{(1,1,1)}$ distinguishes the following pairs of links:*

$L11n325\{1,1\}$	$L11n424\{0,0\}$
$L11n356\{1,0\}$	$L11n434\{0,0\}$
$L10n79\{1,1\}$	$L10n95\{1,0\}$
$L11a404\{1,1\}$	$L11a428\{0,1\}$
$L10n76\{1,1\}$	$L11n425\{1,0\}$
$L11n358\{1,1\}$	$L11n418\{1,0\}$

Proof. These links have three components. The values of F are different on the colored links, obtained by coloring each of these links with three different colors. In Appendix 2, the values of F^3 are listed for the pairs, except for the pair ($L10n76\{1,1\}$, $L11n425\{1,0\}$), for which the invariants are calculated in Section 4 and are shown in Appendix 4.

□

Proposition 3. *The invariant $\mathbf{F}^2_{(2,1)}$ distinguishes the following pairs of links, which are not distinguished by $\mathbf{F}^3_{(1,1,1)}$:*

$L11n358\{0,1\}$	$L11n418\{0,0\}$
$L11a467\{0,1\}$	$L11a527\{0,0\}$



FIGURE 4. Colored links for the calculation of $\mathbf{F}^2_{(2,1)}$

Proof. For every one of the above pairs, we proceed in this way: we firstly calculate F^3 on the links of the pair (all links have three components). These values coincide in each pair, so we obtain that $\mathbf{F}^3_{(1,1,1)}$ does not distinguish these links. Then we calculate $\mathbf{F}^2_{(2,1)}$.

We show here the calculation of $\mathbf{F}^2_{(2,1)}$ on the links of the pair (K, H).

There are three non-equivalent colorations of K and H corresponding to the partitions of type $(2,1)$ of the set of components. The non c-isotopic colored links obtained by such colorations are called respectively $\mathsf{KA}, \mathsf{KB}, \mathsf{KC}$, and $\mathsf{HA}, \mathsf{HB}, \mathsf{HC}$, see Figure 4.



FIGURE 5. Calculation of $F^3(\mathsf{K})$



FIGURE 6. Calculation of $F^3(\mathsf{H})$

From Figures 5 and 6 we obtain the values $F^3(\mathsf{K}) = F^3(\mathsf{H})$, see Appendix 3.

We calculate now $\mathbf{F}^2_{(2,1)}(\mathsf{K}) = \{F(\mathsf{KA}), F(\mathsf{KB}), F(\mathsf{KC})\}$, see Figure 7. Then we compare this set of values with $\mathbf{F}^2_{(2,1)}(\mathsf{H}) = \{F(\mathsf{HA}), F(\mathsf{HB}), F(\mathsf{HC})\}$, obtained from Figure 8. These sets of values, shown in Appendix 3, are different.

□

4. IDENTITIES

For a link with three components, one could define, instead of $\mathbf{F}^2_{(2,1)}$, the invariant defined as the *sum* of the values belonging to the set $\mathbf{F}^2_{(2,1)}$. Surprisingly, despite the fact that the values are all different for the pairs of Proposition 3, their sums coincide. We have indeed:

$$F(\mathsf{KA}) + F(\mathsf{KB}) + F(\mathsf{KC}) - F(\mathsf{HA}) - F(\mathsf{HB}) - F(\mathsf{HC}) = 0$$

$$F(\mathsf{GA}) + F(\mathsf{GB}) + F(\mathsf{GC}) - F(\mathsf{QA}) - F(\mathsf{QB}) - F(\mathsf{QC}) = 0$$

Furthermore, I have found that, in the pairs of Proposition 3, the links are distinguished not only by $\mathbf{F}^3_{1,1,1}$, but by $\mathbf{F}^2_{2,1}$ as well. Naming X and Y the colored links obtained from a pair by coloring the three component with three different colors, and XA, XB, XC, YA, YB, YC, the colored links obtained from that pair by coloring the components with two different colors, the corresponding values of F satisfy:

$$F(\mathsf{XA}) + F(\mathsf{XB}) + F(\mathsf{XC}) - F(\mathsf{YA}) - F(\mathsf{YB}) - F(\mathsf{YC}) = F(\mathsf{X}) - F(\mathsf{Y}).$$

Therefore, in the case of three components, we formulate the following conjecture. Observe that in all the pairs of links here considered, the homflypt polynomial, and then $\mathbf{F}^1_{(3)}$, coincide. Denote by Σ^c the sum of the elements of the set $\mathbf{F}^c_p$.

Conjecture 1. *Let L_1 and L_2 be two links with three components. If $\Sigma^1(L_1) = \Sigma^1(L_2)$, then*

$$\Sigma^3(L_1) - \Sigma^2(L_1) = \Sigma^3(L_2) - \Sigma^2(L_2).$$

The reasons of the above identities are not evident to me and deserve further investigations. Also, the invariants $\mathbf{F}^3_p$ have to be calculated on all pairs of non isotopic links with the same value of F^1.

Example. In Figures 9, 10 and 11 the trees for the calculations of $\mathbf{F}^3_{(1,1,1)}$ and $\mathbf{F}^2_{(2,1)}$ on the links $\mathsf{R} = L10n76\{1,1\}$ and $\mathsf{S} = L11n425\{1,0\}$ are shown. Observe that the values of F on some leaves of the skein-trees for the colored links obtained from R and S, are obtained by using Remark 3, e.g. $\mathsf{R1} = \mathsf{L39}/(wz)$, $\mathsf{R2} = \mathsf{L35}/(wx)$. Also, Remark 5 is used to calculate some values of F, e.g. $F(\mathsf{S4}) = F(\mathsf{L39})F(\mathsf{L5})$, $F(\mathsf{SB5}) = F(\mathsf{L35})F(\mathsf{L4})$.

The values of $\mathbf{F}^3$ and $\mathbf{F}^2_{(2,1)}$ are shown in Appendix 4.

Acknowledgement. A particular thank to Konstantinos Karvounis for having providing me with the list of pairs of non isotopic links that are not distinguished by the homflypt polynomial.



FIGURE 7. Calculation of $F(\mathsf{KA})$, $F(\mathsf{KB})$ and $F(\mathsf{KC})$



FIGURE 8. Calculation of $F(\mathsf{HA})$, $F(\mathsf{HB})$ and $F(\mathsf{HC})$



FIGURE 9. Calculation of F^3 on the links R and S



FIGURE 10. Calculation of $\mathbf{F}^2_{(2,1)}$ on the link R



FIGURE 11. Calculation of $\mathbf{F}^2_{(2,1)}$ on the link S

Appendix 1: List of values of F on simple colored links



Figure 12. Table of simple colored links

	F		F
L1	1	L2	$\frac{tw^2-1}{(1-t)w}$
L3	$\frac{1}{wx}$	L4	$\frac{w(t^2w^2+t-t^2-1)}{1-t}$
L5	$\frac{(1-xtw^2+xt)w}{x}$	L6	$\frac{tw^2+1-t^2w^2-w^2}{tw^3(t-1)}$
L7	$\frac{t-x+xw^2}{tw^3x}$	L8	$\frac{(tw^2-1)^2}{(1-t)^2w^2}$
L9	$\frac{tw^2-1}{(1-t)w^2x}$	L10	$\frac{(t^2w^2+t-t^2-1)(tw^2-1)}{(t-1)^2}$
L11	$\frac{t^2w^2+t-t^2-1}{(1-t)x}$	L12	$\frac{(1-xtw^2+xt)(tw^2-1)}{x(1-t)}$
L13	$\frac{(tw^2+1-t^2w^2-w^2)(1-tw^2)}{tw^4(t-1)^2}$	L14	$\frac{tw^2+1-t^2w^2-w^2}{tw^4(t-1)x}$
L15	$\frac{(t-x+xw^2)(tw^2-1)}{tw^4x(1-t)}$	L16	$\frac{w^2(t^2w^2+t-t^2-1)^2}{(t-1)^2}$
L17	$\frac{w^2(xtw^2-xt-1)(t^2w^2+t-t^2-1)}{x(t-1)}$	L18	$\frac{w^2(1-tw^2+xt^2w^4-2xtw^2+2xt+xt^3w^4-2xt^3w^2-xt^2+xt^3)}{x(t-1)}$
L19	$\frac{(-tw^2-1+t^2w^2+w^2)^2}{(t^2w^6(t-1)^2}$	L20	$\frac{(t-x+xw^2)(tw^2+1-t^2w^2-w^2)}{t^2w^6x(t-1)}$
L21	$\frac{t^3w^2-t^2-2xt^2w^2+xt+2xt^2w^4+x-xw^4t-2xw^2+xw^4}{t^2w^6x(1-t)}$	L22	$\frac{(-tw^2-1+t^2w^2+w^2)(t^2w^2+t-t^2-1)}{(t-1)^2w^2t}$
L23	$\frac{(t-x+xw^2)(t^2w^2+t-t^2-1)}{w^2x(1-t)t}$	L24	$\frac{(xtw^2-xt-1)(-tw^2-1+t^2w^2+w^2)}{w^2x(t-1)t}$
L25	$\frac{-t^2w^2+t-xtw^2-x+xw^2-xt^2w^4+xt^3w^4+xt^2w^2+xt-xt^3w^2}{w^2x(t-1)t}$	L26	$\frac{1}{w^2x^2}$
L27	$\frac{1-xtw^2+xt}{x^2}$	L28	$\frac{t-x+xw^2}{tw^4x^2}$
L29	$\frac{(xtw^2-xt-1)^2*w^2}{x^2}$	L30	$\frac{(t-x+xw^2)^2}{x^2t^2w^6}$
L31	$\frac{(1-xtw^2+xt)(t-x+xw^2)}{w^2tx^2}$	L32	$\frac{w^3(t^3w^2+t^2-t^3-t-t^4w^2+t^4-t^2w^2+1)}{t-1}$
L33	$\frac{-w(t^3w^4+t^2w^2-t^3w^2-tw^2-t^2+2t-1)}{t-1}$	L34	$\frac{-(t^2w^2+t-t^3w^2-tw^2-t^2+t^4w^2-1+w^2)}{t^3w^5(t-1)}$
L35	$\frac{tw^2+1-t^2w^2-w^2-t^3w^4+2t^2w^4-tw^4}{t^2w^5(t-1)}$	L36	$\frac{w^3(1-xtw^2+xt-xt^3w^2+xt^3)}{x}$
L37	$\frac{w(w^2-xw^4t-xt^2w^4+xt^2w^2+xw^2+xt-x)}{x}$	L38	$\frac{t^3-xt^2+xt^2w^2-x+xw^2}{xw^5t^3}$
L39	$\frac{t^2-xt-x+xt^2w^2+xw^2-xt^2w^4+xw^4t}{xw^5t^2}$	L40	$t^2w^2+w^2-t^2w^4$
L41	$\frac{t^2w^2+w^2-1}{t^2w^4}$		

Appendix 2

$F^3(L11n325\{1,1\}) =$
$(-xt^6w^4 + x^2t^6w^6 + xt^5w^4 - x^2t^4 - xtw^2 - 3x^2tw^2 + 6x^2t^2w^2 - 8x^2t^3w^2 + 6x^2t^4w^2 - 3x^2t^5w^2 +$
$x^2t^6w^2 + xt^3w^4 + xt^2w^2 - 2xt^3w^2 - 2x^2t^2w^4 + 5x^2t^3w^4 - 6x^2t^4w^4 + 5x^2t^5w^4 - 2x^2t^6w^4 - x^2 +$
$xt + xt^4w^2 - xt^5w^2 + xt^6w^2 - xt^2 - 4x^2t^2 + 3x^2t - 2x^2t^5w^6 + x^2t^4w^6 + x^2w^2 + 3x^2t^3 + t^5w^2 +$
$xt^3 - xt^4)/(w^6x^2t^5)$
$F^3(L11n424\{0,0\}) =$
$-(2xt^6w^4 - 2xt^5w^4 + x^2t^4 + 2x^2tw^2 - 7x^2t^2w^2 + 8x^2t^3w^2 - 5x^2t^4w^2 + 4x^2t^5w^2 - x^2t^6w^2 +$
$2xt^3w^2 + 3x^2t^2w^4 - 4x^2t^3w^4 + 6x^2t^4w^4 - 6x^2t^5w^4 + x^2t^6w^4 + x^2 - xt^4w^2 + 2xt^5w^2 - 2xt^6w^2 +$
$4x^2t^2 - 2x^2t - xt^4w^4 + 2x^2t^5w^6 - 2x^2t^4w^6 - x^2w^2 - 4x^2t^3 - t^5w^2 - 2xt^3 + 2xt^4)/(w^6x^2t^5)$

$F^3(L11n356\{1,0\}) =$
$(-t^2x^2 + 2tx^2 - xt^5w^4 + xt^3w^2 - x^2 + 8t^3x^2w^4 - x^2t^6w^4 + t^2w^2 + x^2t^6w^6 - 4t^2x^2w^4 - xt^3w^4 -$
$5t^3x^2w^2 - 3x^2t^5w^6 - 4x^2t^4w^4 + x^2t^4w^2 + 3x^2t^5w^4 + 2x^2w^2 + xt^5w^2 + 5x^2t^2w^2 - 5x^2tw^2 -$
$x^2w^4 - xtw^2 + xtw^4 - 3x^2t^3w^6 + 3tx^2w^4 + 3x^2t^4w^6)/(w^2x^2t^2)$
$F^3(L11n434\{0,0\}) =$
$(xt^3 - 2t^2x^2 + 2tx^2 - xt + x + xt^3w^2 + 8t^3x^2w^4 + xt^4w^2 - x^2t^6w^4 - xt^4w^4 + t^2w^2 + x^2t^6w^6 -$
$3t^2x^2w^4 - 2xt^3w^4 - 4t^3x^2w^2 - 2x^2t^5w^6 - 5x^2t^4w^4 + 2x^2t^4w^2 + 2x^2t^5w^4 + x^2w^2 - xt^2 +$
$5x^2t^2w^2 - xw^2 - 6x^2tw^2 - x^2w^4 + xtw^4 - 4x^2t^3w^6 + xt^2w^4 + 4tx^2w^4 + 3x^2t^4w^6)/(w^2x^2t^2)$

$F^3(L10n79\{1,1\}) =$
$-(-2x^2t^2w^4 + x^2t^3w^4 + 3t^4w^2x^2 - 2x^2t^4w^4 + 2x^2t^3 - 3w^2x^2t^3 + t^7x^2w^4 - x^2t^4 + x^2t^6 +$
$t^7xw^2 + x^2t^5 - 2t^5xw^2 + 2t^5x - t^7x - x^2tw^2 - x^2t^7w^2 + x^2w^2 - 3x^2t^5w^2 - x^2 - t^6 + 2x^2t^5w^4 +$
$5w^2x^2t^2 - 2xt^3w^2 + 2t^3x - 3x^2t^2 + x^2t - x^2t^6w^2)/(w^8x^2t^6)$
$F^3(L10n95\{1,0\}) =$
$(t^6 - 3w^2x^2t^2 + x^2t^2w^4 - 2x^2t + 2x^2tw^2 - 2x^2t^5w^4 + 2x^2t^2 + x^2t^4 + t^5xw^2 - 2x^2t^3 + xt^3w^2 -$
$t^3x - t^5x + 2x^2t^4w^4 + 2x^2t^5w^2 + 4w^2x^2t^3 + x^2t^6w^4 - 2x^2t^3w^4 - x^2w^2 - x^2t^6w^2 + xtw^2 -$
$xt + x^2 - 3t^4w^2x^2)/(w^8x^2t^6)$

$F^3(L11a404\{1,1\}) =$
$-(xt^2w^4 - xw^2 + x - 2xt^6w^4 - 2x^2t^7w^2 + x^2t^8w^2 + 4x^2t^7w^4 - 2x^2t^8w^4 + 2x^2t^6w^6 - 2x^2t^7w^6 +$
$x^2t^8w^6 - 4x^2t^4 + xtw^2 - 3x^2tw^2 + 4x^2t^2w^2 - 15x^2t^3w^2 + 9x^2t^4w^2 - 15x^2t^5w^2 + 6x^2t^6w^2 -$
$xt^3w^4 - 4xt^2w^2 + xt^3w^2 - t^5 - x^2t^2w^4 + 8x^2t^3w^4 - 6x^2t^4w^4 + 15x^2t^5w^4 - 7x^2t^6w^4 -$
$xt - xt^4w^2 + 4xt^6w^2 - t^3 + 3xt^2 - 3x^2t^2 + 2x^2t + 2xt^4w^4 - x^2t^3w^6 + x^2tw^4 - 4x^2t^5w^6 +$
$x^2t^4w^6 + 8x^2t^3 + t^5w^2 + 4x^2t^5 - x^2t^6 - 2xt^6 - xt^4)/(x^2t^3)$
$F^3(L11a428\{0,1\}) =$
$-(-t^6x - 3x^2t^4 + t^4x + 4x^2t^5 + 7x^2t^3 - t^3 + t^5xw^2 - x^2 + t^5w^2 - 4x^2t^2 + 2t^2x + 9x^2t^4w^2 + 2x^2t +$
$x^2t^7 + 6x^2t^3w^4 - t^8x + x^2w^2 - t^5 - x^2t^8w^4 - 2x^2t^2w^4 - 7x^2t^4w^4 - 16x^2t^5w^2 - 3x^2t^7w^6 +$
$4x^2t^6w^2 + 2x^2t^6w^6 + 6x^2t^2w^2 + 2t^6xw^2 - xt^6w^4 + x^2t^8w^6 + x^2t^4w^6 - 6x^2t^6w^4 + 2xt^4w^4 -$
$13x^2t^3w^2 + 16x^2t^5w^4 - 4x^2t^7w^2 - 4x^2t^5w^6 - t^5x + 6x^2t^7w^4 - 3t^4xw^2 + t^8xw^2 - t^7xw^4 -$
$2x^2tw^2 + t^7xw^2 - 2xt^2w^2)/(x^2t^3)$

$F^3(L11n358\{1,1\}) =$
$(-x^2t + x^2t^2 - x^2t^3 + x^2t^4 + 2x^2tw^2 - 2x^2t^2w^2 + 2x^2t^2w^4 - x^2tw^4 + 3x^2t^3w^2 - 3x^2t^4w^2 -$
$4x^2t^3w^4 + 2x^2t^3w^6 + 2x^2t^4w^4 - x^2t^2w^6 - 4x^2t^5w^4 + x^2t^4w^8 + x^2t^5w^6 + 3x^2t^5w^2 -$
$x^2t^4w^6 - t^6w^8x^2 + 2t^6w^4x^2 - t^6w^2x^2 + t^7w^6x^2 - t^7w^4x^2 - x + tw^2 + 2xt^3w^4 + xt -$
$2xtw^2 + xw^2 + 4xt^2w^2 - 4xt^2w^4 - xt^3w^2 + xw^4t + xw^2t^4 - w^6xt^3 + xw^6t^4 - 2xw^4t^4)w^2/(x^2t)$
$F^3(L11n418\{1,0\}) =$
$(-2x^2t + x^2t^2 - x^2t^3 + x^2t^4 + 3x^2tw^2 - 2x^2t^2w^2 + 2x^2t^2w^4 + 4x^2t^3w^2 - 3x^2t^4w^2 - 5x^2t^3w^4 +$
$x^2t^3w^6 + x^2t^4w^4 - x^2t^2w^6 - 4x^2t^5w^4 + x^2t^5w^6 + 3x^2t^5w^2 + x^2t^4w^6 + x^2 - t^6w^8x^2$
$+2t^6w^4x^2 - t^6w^2x^2 - 2x^2w^2 + t^7w^6x^2 - t^7w^4x^2 + w^4x^2 - w^6xt^2 + tw^2 - 2xt^3w^4 - xt + 2xtw^2 -$
$xw^2 + xt^2 + xt^2w^2 - xt^2w^4 + xt^3w^2 + xw^4 - xw^4t + w^6xt^3 + w^8x^2t^3 - w^6x^2t)w^2/(x^2t)$

Appendix 3

$$\mathbf{F}^3_{(1,1,1)}(L11n358\{0,1\}) = \mathbf{F}^3_{(1,1,1)}(L11n418\{0,0\})$$

$F(\mathsf{K}) = F(\mathsf{H}) = (-tx^2w^4 + x^2t^5w^2 + xt^2w^2 + t^3x^2 + t^5 - 3t^4x - xw^4t^6 - t^2w^6x^2 - t^4x^2 + t^5x + x^2t^7w^6 - x^2t^7w^4 + xt^6w^2 + xw^4t^7 - xt^7w^2 + t^2x^2 - 2x^2w^2 - xt^2 + x^2w^4 + 2x^2t^3w^6 + x^2t^6w^2 - 3x^2t^4w^6 + x^2tw^2 - 5x^2t^2w^2 + 5x^2t^4w^4 + 2x^2t^5w^6 + x^2 + t^3x^2w^2 + 5t^2x^2w^4 - 4t^3x^2w^4 - x^2t^6w^6 + 3xt^4w^2 - x^2t^4w^2 - xt^5w^2 - 3x^2t^5w^4)/(t^5x^2w^8)$

$$\mathbf{F}^2_{(2,1)}(L11n358\{0,1\}) \neq \mathbf{F}^2_{(2,1)}(L11n418\{0,0\}))$$

$\mathbf{F}^2_{(2,1)}(L11a435\{0,0\}) = \{F(\mathsf{KA}), F(\mathsf{KB}), F(\mathsf{KC})\}$

$F(\mathsf{KA}) = (2t^3w^2 + t^4w^4 - 4t^4w^2 - 2t^3 + 6xt^2w^2 - 2t^5 + 5t^5w^2 + t^4x - 2t^5w^4 + 2t^6w^4 - 4t^6w^2 + 3t^4 - 3w^6xt^3 + 5w^6xt^4 + w^6xt^2 - 6xw^6t^5 + 4xw^6t^6 - 4xw^4t^6 - t^5x + 2t^7w^2 - t^8w^2 + t^8w^4 - 2t^7w^4 + xw^4t^7 + xt^7w^2 - 2xt^7w^6 - t^2w^2 + xt - xt^3 - xt^2 + 2xw^2 - x + t^6 + 7xt^5w^4 + t^2 - 5xt^3w^2 - xw^4 + 3xt^4w^2 + 9xt^3w^4 - 3xtw^2 + 2xtw^4 - 9xt^4w^4 - 6xt^2w^4 + xt^8w^6 - xt^8w^4)/((t-1)w^8t^5x)$,

$F(\mathsf{KB}) = -(-7xt^2w^2 - t^5 + t^4x + t^6w^2 + 3w^6xt^3 - 5w^6xt^4 - w^6xt^2 + 5xw^6t^5 - 3xw^6t^6 + 3xw^4t^6 - xw^4t^7 + xt^7w^6 - xt + 2xt^2 - 2xw^2 + x - 5xt^5w^4 + 5xt^3w^2 + xw^4 - 5xt^4w^2 - 8xt^3w^4 + 3xtw^2 - 2xtw^4 + 9xt^4w^4 + 6xt^2w^4)/((t-1)w^8t^5x)$,

$F(\mathsf{KC}) = -(t^3w^2 + t^4w^4 - t^4w^2 - t^3 - 7xt^2w^2 + t^5w^2 - 2t^5w^4 + t^6w^4 + 3w^6xt^3 - 5w^6xt^4 - w^6xt^2 + 4xw^6t^5 - 2xw^6t^6 + 2xw^4t^6 - xw^4t^7 + xt^7w^6 - xt - xt^3 + 2xt^2 - 2xw^2 + x - 6xt^5w^4 + 6xt^3w^2 + xw^4 - 4xt^4w^2 - 8xt^3w^4 + 2xt^5w^2 + 3xtw^2 - 2xtw^4 + 9xt^4w^4 + 6xt^2w^4)/((t-1)w^8t^5x)$.

$\mathbf{F}^2_{(2,1)}(L11n418\{0,0\}) = \{F(\mathsf{HA}), F(\mathsf{HB}), F(\mathsf{HC})\}$

$F(\mathsf{HA}) = (-t^4w^2 + 7xt^2w^2 - t^5 + 2t^5w^2 + t^6w^4 - 3t^6w^2 + t^4 - 3w^6xt^3 + 5w^6xt^4 + w^6xt^2 - 5xw^6t^5 + 3xw^6t^6 - 3xw^4t^6 - t^5x + 2t^7w^2 - t^8w^2 + t^8w^4 - 2t^7w^4 + xw^4t^7 + xt^7w^2 - 2xt^7w^6 + xt - 2xt^2 + 2xw^2 - x + t^6 + 6xt^5w^4 - 5xt^3w^2 - xw^4 + 4xt^4w^2 + 8xt^3w^4 - 3xtw^2 + 2xtw^4 - 9xt^4w^4 - 6xt^2w^4 + xt^8w^6 - xt^8w^4)/(w^8t^5x(t-1))$,

$F(\mathsf{HB}) = -(-t^3w^2 + t^4w^2 + t^3 - 6xt^2w^2 - t^5w^2 + t^6w^2 - t^4 + 3w^6xt^3 - 5w^6xt^4 - w^6xt^2 + 5xw^6t^5 - 3xw^6t^6 + 3xw^4t^6 - xw^4t^7 + xt^7w^6 + t^2w^2 - xt + xt^2 - 2xw^2 + x - 6xt^5w^4 - t^2 + 6xt^3w^2 + xw^4 - 4xt^4w^2 - 9xt^3w^4 + xt^5w^2 + 3xtw^2 - 2xtw^4 + 9xt^4w^4 + 6xt^2w^4)/(w^8t^5x(t-1))$,

$F(\mathsf{HC}) = -(t^4w^2 - 7xt^2w^2 - t^5w^2 + t^6w^2 - t^4 + 3w^6xt^3 - 5w^6xt^4 - w^6xt^2 + 5xw^6t^5 - 3xw^6t^6 + 3xw^4t^6 - xw^4t^7 + xt^7w^6 - xt + 2xt^2 - 2xw^2 + x - 6xt^5w^4 + 5xt^3w^2 + xw^4 - 4xt^4w^2 - 8xt^3w^4 + xt^5w^2 + 3xtw^2 - 2xtw^4 + 9xt^4w^4 + 6xt^2w^4)/(w^8t^5x(t-1))$.

$$\mathbf{F}^3_{(1,1,1)}(L11a467\{0,1\}) = \mathbf{F}^3_{(1,1,1)}(L11a527\{0,0\})$$

$F(\mathsf{G}) = F(\mathsf{Q}) = -(3xt^4 - x^2t^2 - x^2t^3 + 3x^2w^2 + 9x^2t^2w^2 + 14x^2t^3w^4 + 5x^2tw^4 - 10x^2t^3w^6 - 4x^2t^3w^2 - 3xt^4w^2 - 13x^2t^4w^4 + 13x^2t^4w^6 + 2x^2t^4w^2 + w^8x^2t^3 - 3t^4w^8x^2 + t^7x^2w^4 - t^6x^2w^2 - 2t^7w^6x^2 + xt^7w^2 - t^5 + 7t^5x^2w^4 - 10t^5x^2w^6 - t^6x^2w^4 + 5t^6x^2w^6 + t^5xw^2 + t^6w^4x - t^7xw^4 - t^6xw^2 + 4w^8t^5x^2 - x^2w^6t - t^5x^2w^2 + t^7w^8x^2 - 3t^6w^8x^2 + x^2t^4 - xt^5 - x^2 - xt^2w^2 + xt^2 + 5w^6x^2t^2 - 4w^2tx^2 - 2x^2w^4 - 13w^4x^2t^2)/(w^8t^5x^2)$

$$\mathbf{F}^2_{(2,1)}(L11a467\{0,1\}) \neq \mathbf{F}^2_{(2,1)}(L11a527\{0,0\})$$

$\mathbf{F}^2_{(2,1)}(L11a467\{0,1\}) = \{F(\mathsf{GA}), F(\mathsf{GB}), F(\mathsf{GC})\}$

$F(\mathsf{GA}) = (-8xt^4w^2 + 27xt^4w^4 - 23xt^4w^6 + 15t^3w^6x - 6t^2w^6x - 7w^8t^5x - 18t^5xw^4 + 22t^5w^6x + t^4w^4 - 2t^5w^4 + xw^6t + 3t^5xw^2 - 4t^7w^8x + 7t^6w^8x + 4t^4w^8x - 14t^6w^6x + 7t^6w^4x - 2t^7xw^4 - t^3w^8x + t^5w^2 + t^6w^4 + x - 26xt^3w^4 - 3xw^2 + 2xw^4 - xt^8w^6 + xt^8w^8 + 7xtw^2 + 6t^7w^6x - xt - 14xt^2w^2 + 2xt^2 + t^3w^2 - xt^3 + 18xt^2w^4 + 13xt^3w^2 - 7xw^4t - t^4w^2 - t^3)/(xt^5w^8(1-t))$,

$F(\mathsf{GB}) = (-xt^4 - 7xt^4w^2 + 27xt^4w^4 - 23xt^4w^6 + 15t^3w^6x - 6t^2w^6x - xt^7w^2 - 7w^8t^5x + 2t^5 - 19t^5xw^4 + 24t^5w^6x - t^4w^4 + 2t^5w^4 + xw^6t + t^5xw^2 - 4t^7w^8x + 7t^6w^8x + 4t^4w^8x - 16t^6w^6x + 9t^6w^4x - 2t^7xw^4 - t^3w^8x - 5t^5w^2 + 2t^7w^4 - 2t^7w^2 - 2t^6w^4 + 4t^6w^2 + xt^5 - t^8w^4 - t^6 + t^8xw^4 + x - 27xt^3w^4 - 3xw^2 + t^2w^2 - t^2 + 2xw^4 - 2xt^8w^6 + xt^8w^8 + 7xtw^2 + 7t^7w^6x - xt - 13xt^2w^2 + xt^2 - 2t^3w^2 + xt^3 + 18xt^2w^4 + 12xt^3w^2 - 7xw^4t + 4t^4w^2 + t^8w^2 - 3t^4 + 2t^3)/(xt^5w^8(1-t))$,

$F(\mathsf{GC}) = (xt^4 - 9xt^4w^2 + 27xt^4w^4 - 23xt^4w^6 + 15t^3w^6x - 6t^2w^6x - 7w^8t^5x - t^5 - 17t^5xw^4 + 23t^5w^6x + xw^6t + t^5xw^2 - 4t^7w^8x + 7t^6w^8x + 4t^4w^8x - 15t^6w^6x + 8t^6w^4x - 2t^7xw^4 - t^3w^8x + t^6w^2 + x - 26xt^3w^4 - 3xw^2 + 2xw^4 - xt^8w^6 + xt^8w^8 + 7xtw^2 + 6t^7w^6x - xt - 14xt^2w^2 + 2xt^2 + 18xt^2w^4 + 12xt^3w^2 - 7xw^4t)/(xt^5w^8(1-t))$.

$\mathbf{F}^2_{(2,1)}(L11a527\{0,0\}) = \{F(\mathsf{QA}), F(\mathsf{QB}), F(\mathsf{QC})\}$

$F(\mathsf{QA}) = (-8xt^4w^2 + 27xt^4w^4 - 23xt^4w^6 + 15t^3w^6x - 6t^2w^6x - 7w^8t^5x - 18t^5xw^4 + 23t^5w^6x + xw^6t + 2t^5xw^2 - 4t^7w^8x + 7t^6w^8x + 4t^4w^8x - 15t^6w^6x + 8t^6w^4x - 2t^7xw^4 - t^3w^8x - t^5w^2 + t^6w^2 + x - 26xt^3w^4 - 3xw^2 + 2xw^4 - xt^8w^6 + xt^8w^8 + 7xtw^2 + 6t^7w^6x - xt - 14xt^2w^2 + 2xt^2 + 18xt^2w^4 + 12xt^3w^2 - 7xw^4t + t^4w^2 - t^4)/(xt^5w^8(1-t))$,

$F(\mathsf{QB}) = (-8xt^4w^2 + 27xt^4w^4 - 23xt^4w^6 + 15t^3w^6x - 6t^2w^6x - xt^7w^2 - 7w^8t^5x + t^5 - 18t^5xw^4 + 23t^5w^6x + xw^6t + t^5xw^2 - 4t^7w^8x + 7t^6w^8x + 4t^4w^8x - 15t^6w^6x + 8t^6w^4x - 2t^7xw^4 - t^3w^8x - 2t^5w^2 + 2t^7w^4 - 2t^7w^2 - t^6w^4 + 3t^6w^2 + xt^5 - t^8w^4 - t^6 + t^8xw^4 + x - 26xt^3w^4 - 3xw^2 + 2xw^4 - 2xt^8w^6 + xt^8w^8 + 7xtw^2 + 7t^7w^6x - xt - 14xt^2w^2 + 2xt^2 + 18xt^2w^4 + 12xt^3w^2 - 7xw^4t + t^4w^2 + t^8w^2 - t^4)/(xt^5w^8(1-t))$,

$F(\mathsf{QC}) = (-8xt^4w^2 + 27xt^4w^4 - 23xt^4w^6 + 15t^3w^6x - 6t^2w^6x - 7w^8t^5x - 18t^5xw^4 + 23t^5w^6x + xw^6t + 2t^5xw^2 - 4t^7w^8x + 7t^6w^8x + 4t^4w^8x - 15t^6w^6x + 8t^6w^4x - 2t^7xw^4 - t^3w^8x - t^5w^2 + t^6w^2 + x - 27xt^3w^4 - 3xw^2 + t^2w^2 - t^2 + 2xw^4 - xt^8w^6 + xt^8w^8 + 7xtw^2 + 6t^7w^6x - xt - 13xt^2w^2 + xt^2 - t^3w^2 + 18xt^2w^4 + 13xt^3w^2 - 7xw^4t + t^4w^2 - t^4 + t^3)/(xt^5w^8(1-t))$.

Appendix 4

$\mathbf{F}^3_{(1,1,1)}$ and $\mathbf{F}^2_{(2,1)}$ on the links $\mathsf{R} = L10n76\{1,1\}$ and $\mathsf{S} = L11n425\{1,0\}$.

$\mathbf{F}^3_{(1,1,1)}(\mathsf{R}) = \{F^3(\mathsf{R})\}$

$F^3(\mathsf{R}) = (-x^2w^4t+t^3w^6x^2+3x^2t^2w^4-4x^2t^3w^4+t^5xw^2-t^5xw^4-x^2t^4w^2+3x^2t^3w^2-x^2t^5w^4-4x^2t^2w^2+x^2t^5w^6+3x^2t^4w^4-2x^2t^4w^6+t^5-t^4x+x^2t^2-2x^2t-x^2w^2+x^2+3w^2tx^2-xt+xtw^2-xt^2w^2+t^4w^4x+xt^2w^4-xt^3w^4+2xt^3w^2-xt^3)/(t^5x^2w^8)$.

$\mathbf{F}^2_{(2,1)}(\mathsf{R}) = \{F(\mathsf{RA}), F(\mathsf{RB}), F(\mathsf{RC})\}$

$F(\mathsf{RA}) = (-3t^5w^6x-t^5xw^2+4t^5xw^4+4xt^4w^2+2t^5w^4-t^5w^2-t^6w^4+t^3+t^6w^6x-t^6w^4x-tw^2+t-x+2t^4w^2+3xt-4xtw^2+xw^2-t^2w^4+2t^3w^4+2t^2w^2-t^2+7xt^2w^2-3xt^2-7t^4w^4x-t^3w^6x+3t^4w^6x-2t^4w^4-4xt^2w^4+7xt^3w^4-7xt^3w^2-3t^3w^2+xt^3+xw^4t)/(w^8t^5x(t-1))$,

$F(\mathsf{RB}) = (-4t^5w^6x+t^5xw^2+3t^5xw^4+5xt^4w^2+t^5-t^4x-t^6w^2+2t^6w^6x-2t^6w^4x-x+2xt-3xtw^2+xw^2+7xt^2w^2-3xt^2-8t^4w^4x-2t^3w^6x+4t^4w^6x-4xt^2w^4+7xt^3w^4-5xt^3w^2+xw^4t)/(w^8t^5x(t-1))$,

$F(\mathsf{RC}) = (-4t^5w^6x+t^5xw^2+3t^5xw^4+5xt^4w^2+t^5-t^4x-t^6w^2+2t^6w^6x-2t^6w^4x-x+2xt-3xtw^2+xw^2+7xt^2w^2-3xt^2-8t^4w^4x-2t^3w^6x+4t^4w^6x-4xt^2w^4+7xt^3w^4-5xt^3w^2+xw^4t)/(w^8t^5x(t-1))$.

$\mathbf{F}^3_{(1,1,1)}(\mathsf{S}) = \{F^3(\mathsf{S})\}$

$F^3(\mathsf{S}) = -(x^2w^4t-2t^3w^6x^2-3x^2t^2w^4+4x^2t^3w^4-2t^5xw^2+2t^5xw^4+x^2t^4w^2-2x^2t^3w^2+x^2t^5w^4+5x^2t^2w^2-4x^2t^4w^4+2x^2t^4w^6-t^5+2t^4x-2x^2t^2+x^2t+x^2w^2-x^2-t^6x+t^5x^2+t^4x^2-t^6x^2w^2-2t^5x^2w^2+t^6x^2w^4+t^6xw^2-2w^2tx^2-2t^4w^4x-2xt^3w^2+2xt^3)/(t^5x^2w^8)$.

$\mathbf{F}^2_{(2,1)}(\mathsf{S}) = \{F(\mathsf{SA}), F(\mathsf{SB}), F(\mathsf{SC})\}$

$F(\mathsf{SA}) = (-3t^5w^6x-t^5xw^2+4t^5xw^4+4xt^4w^2+2t^5w^4-t^5w^2-t^6w^4+t^3+t^6w^6x-t^6w^4x-x+t^4w^2+2xt-3xtw^2+xw^2+7xt^2w^2-3xt^2-8t^4w^4x-2t^3w^6x+4t^4w^6x-t^4w^4-4xt^2w^4+7xt^3w^4-6xt^3w^2-t^3w^2+xt^3+xw^4t)/(w^8t^5x(t-1))$,

$F(\mathsf{SB}) = -(4t^5w^6x-t^5xw^2-3t^5xw^4-5xt^4w^2-t^5+t^4x-2t^6w^6x+2t^6w^4x+t^6x+x-t^6xw^2-t^7+t^6+t^7w^2-t^7xw^2+t^7xw^4-2xt+3xtw^2-xw^2-7xt^2w^2+3xt^2+8t^4w^4x+2t^3w^6x-4t^4w^6x+4xt^2w^4-7xt^3w^4+5xt^3w^2-xw^4t)/(w^8t^5x(t-1))$,

$F(\mathsf{SC}) = (-3t^5w^6x-t^5xw^2+4t^5xw^4+4xt^4w^2+2t^5w^4-t^5w^2-t^6w^4+t^3+t^6w^6x-t^6w^4x-x+t^4w^2+2xt-3xtw^2+xw^2+7xt^2w^2-3xt^2-8t^4w^4x-2t^3w^6x+4t^4w^6x-t^4w^4-4xt^2w^4+7xt^3w^4-6xt^3w^2-t^3w^2+xt^3+xw^4t)/(w^8t^5x(t-1))$.

References

[1] *An invariant of colored links via skein relation*, to appear in Arnold Math. Journal.
[2] F. Aicardi, J. Juyumaya, *Tied Links*, to appear in JKTR, Slavik Jablan memorial volume, see also arXiv:1503.00527.
[3] F. Aicardi, J. Juyumaya, *Markov trace on the algebra of braids and ties.* to be published in Moscow Math. J. See also arXiv:1408.5672.
[4] J. C. Cha, C. Livingston, LinkInfo: Table of Knot Invariants, http://www.indiana.edu/∼linkinfo, April 16, 2015.

The Abdus Salam Centre for Theoretical Physics (ICTP), Strada Costiera, 11, 34151 Trieste, Italy.

E-mail address: faicardi@ictp.it